

SCOPIA® GULLIVER™

COMPACT

Sutera cordata (Bacopa)

- Large flower size with good branching
- Early flowering
- Well suited for combinations
- Naturally controlled and well-branched habit



Compact



30-36 cm
12-14 in



10-15 cm
4-6 in



Pollinator
Friendly



Hanging
Basket



12-15 cm
Qt, Gal



SCOPIA® GULLIVER™ Compact series



SCOPIA® GULLIVER™
Compact Rosa



SCOPIA® GULLIVER™
Compact Blush



SCOPIA® GULLIVER™
Compact White




SCOPIA® GULLIVER™
Compact Purple

SCOPIA® GULLIVER™ COMPACT

For additional information and availability:

SCOPIA® GULLIVER™ COMPACT 

Bacopa / Sutera



Stick on priority – Number 1 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 72°/74°F (22°/23°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 65° - 68° (18° - 20° C)		Weeks 3-4 100 to 150 ppm N	Day of sticking & Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6/6.2	0.80	72°/74°F (22°/23°C)	50 ppm N in mist				Day of sticking
Week 2	5.6/6.2	.80 to .90	72°/74°F (22°/23°C)	50 ppm N in mist				Second fungicide app
Week 3	5.6/6.2	.90 to 1.0	Cool to 68°/70°F (20°/21°C)	Feed 100 ppm to 150 ppm				
Week 4	5.6/6.2	1.0 to 1.2	65°/68°F (18°/20°C)	Feed 100 ppm to 150 ppm		Daminozide if needed		



Irrigation specification - Remove from mist as soon as possible to avoid leaf disease issues. Elevated humidity at the table level is better than misting on the foliage.



The Scopia Gulliver™ Compact Series is well branched. Spray fungicide to control leaf diseases.

Bacopa / Sutera

- Maintain lower pH levels for optimal Iron (Fe) and Manganese (Mn) availability to plants
- Maintain soil moisture at medium level to maximize root structure
- Cool temperature is best for high quality finishing

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks	Average Day	Pinch / Optional No lighting necessary	100 - 150 ppm N	Spray application of Daminozide if required
(12 cm) 5" pot 1 ppp	65°-68° F (18°-20° C)		Soil EC 0.8 - 1.0 pH 5.6 to 6.0	
8 to 9 wks	65°-68° F (18°-20° C)			
(15 cm) 6 ½" pot 1 ppp				
8 to 9 wks				
(20 cm) 8" pot 3 ppp				
8 to 9 wks				
(25 cm) 10" pot 3 ppp				

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIACARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRADIONE, POLYOXIN
	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.6 to 6.0	0.8 to 1.0	65° to 68° F (18° to 20° C)	Feed 100 to 150 ppm	Day Neutral	Daminozide spray application if needed	Spray fungicide after transplant	
Week 6 to 12	5.6 to 6.0	0.8 to 1.0	65° to 68° F (18° to 20° C)	Feed 100 to 150 ppm	Not Required			



- Consistent soil moisture levels are best – allow soil to dry slightly between irrigations
- Fungicide applications and good air movement to reduce disease pressures
- Scout for Aphids on regular basis