

DIWSLI SHOER™

Euphorbia hypericifolia

- Low maintenance
- Heat and drought tolerant
- Great filler product, great for combinations



Upright-
Bush



28-35 cm
12-14 in



20-25 cm
8-10 in



Heat
tolerant



Euphorbia hypericifolia



Stick on priority – Number 3 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 72° - 74° F (22° - 23° C)	Recommended	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 65° - 68° (18° - 20° C)		Weeks 3-5 100 to 150 ppm N	Spray fungicide to control Botrytis and bacteria Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 to 6.2	0.80	72° - 74° F (22° - 23° C)	50 ppm N in mist			Day of sticking	Feeding through Mist
Week 2	5.8 to 6.2	.80 to .90	72° - 74° F (22° - 23° C)	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	Cool to 68° - 70° F (20° - 21° C)	Feed 100 ppm to 150 ppm				
Week 4	5.8 to 6.2	1.0 to 1.2	65° - 68° F (18° - 20° C)	Feed 100 ppm to 150 ppm				
Week 5	5.8 to 6.2	1.0 to 1.2	65° - 68° F (18° - 20° C)	Feed 100 ppm to 150 ppm		Daminozide as needed		



- **Irrigation specification** – Spray adjuvant after sticking.



- Application of hormone as dip or K-IBA spray can hasten rooting
- Pinch is not recommended / very sensitive to ethephon applications
- Maintain moderate damp soil during rooting



Euphorbia hypericifolia

- Pinch is not recommended
- Over wet soil in low light conditions can develop root disease issues, allow soil to dry between irrigations

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks (12 cm) 5" pot 1 ppp	Average Day 60° - 68° (16° - 20° C)	Pinch not recommended	150 - 200 ppm N	
7 to 8 wks (15 cm) 6 ½" 1 ppp			Soil EC 1.0 - 1.2 pH 5.8 to 6.2	
7 to 8 wks (20 cm) 8"	3 plants per container	No lighting necessary		
7 to 8 wks (25 cm) 10"	3 plants per container			

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.8 to 6.2	1.0 to 1.2	65° to 68° F (18° to 20° C)	feed 150 - 200 ppm		Daminozide as needed	Spray fungicide after transplant	
Week 6 to 12	5.8 to 6.2	1.0 to 1.2	60° to 68° F (16° to 20° C)	feed 150 - 200 ppm	Not Required	Paclobutrazol Drench at 1 to 2 ppm		Paclobutrazol 1 to 2 ppm drench when plants reach 85% of desired size

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRDIONE, POLYOXIN
	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB

FINISHING TIPS



- Consistent soil moisture levels are best – allow soil to dry slightly between irrigations
- Scout for Aphids, Thrips and Whiteflies