

# MUSICA®

*Impatiens walleriana*

- Spectacular double flowers in a wide range of colors
- Branched, compact growth with uniform habit within the series
- No pinching is necessary
- Superb for pots, hanging baskets and window boxes, under semi-shaded areas




Mounded




Early



Hanging  
Basket



25-35 cm  
10-15 in



35-45 cm  
15-20 in



MUSICA®

DANZIGER

# The MUSICA® Collection



Musica® Pink Aroma



Musica® Lavender

Best Seller



Musica® Bicolor Dark Red



Musica® Bicolor Pink

Best Seller



Musica® Electric Purple



Musica® Elegant Red



## *Impatiens walleriana*



Stick on priority – Number 2 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 72°-74°F (22°-23°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 68° - 70° (20° - 21° C)		Weeks 3-4 100 ppm N	Apply broad spectrum Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 to 6.2	0.80	72°-74°F (22°-23°C)	50 ppm N in mist				Day of sticking
Week 2	5.8 to 6.2	.80 to .90	72°-74°F (22°-23°C)	50 ppm N in mist				Second fungicide app
Week 3	5.8 to 6.2	.80 to .90	Cool to 68° - 70° (20° - 21° C)	Feed 100 ppm N				
Week 4	5.8 to 6.2	.80 to .90	68° - 70° (20° - 21° C)	Feed 100 ppm N				Should not be required



- Spray adjuvant at sticking. Remove from mist as soon as possible



- Shade will be required for later spring/summer production
- Maintain warm temperatures allowing plants to develop
- Average daily temperature of 70° F/ 20° C
- Spray fungicide to control botrytis



## *Impatiens walleriana*

- Shade will be required for later spring/summer production
- Maintain a low EC level for proper growth and sizing of plant
- Transplant liners when plants are rooted and actively growing

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks (12 cm) 5" pot 1 ppp	Average Day 65° to 68° F (18° to 20° C)	Pinch –No  Daylength Neutral	100 – 125 ppm N  Soil EC 0.8 - 1.0 pH 5.8 to 6.2	Paclbutrazol at 0.5 ppm drench within 3 weeks of finishing. Allowing soil to dry between irrigations and modest fertility will keep plants more compact
8 to 9 wks (15 cm) 6 ½" pot 1 ppp	65° to 68° F (18° to 20° C)			

<b>Pests</b>	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrip	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
<b>Diseases</b>	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 5 Transplant</b>	5.8 to 6.2	0.8 to 1.0	65° to 68° F (18° to 20° C)	feed 100 to 125 ppm	Day neutral	Paclbutrazol at 0.5 ppm drench within 3 weeks of finishing	Drench fungicide at transplant	
<b>Week 6 to 13</b>	5.8 to 6.2	0.8 to 1.0	65° to 68° F (18° to 20° C)	feed 100 to 125 ppm				

FINISHING TIPS



- Spray broad spectrum fungicide after transplant
- Scout and spray for Powdery Mildew
- Scout for Aphids, Leaf Miners and Thrips on regular basis