CAPELLA™

Petunia hybrida

- Naturally, controlled habit was selected for quarts /12cm production, will continue to gain size to fill a larger container or garden bed
- Performs well in propagation
- Excellent branching creates a tidy small pot with minimal PGR
- Blooms early enough for any market

Mounded

cm 20-25
8-10 in

cm 25-36
10-14 in

Hanging Basket

12-15 Qt, Patio Pot

Landscape

Ground Cover

Spring / Summer Flowering

The **CAPELLA™** series

CAPELLA™ Inked
PE-22-3891

CAPELLA™ Rose Ink
PE-24-4624

Capella™ Hello Sunshine

Capella™ Berry Lace

Capella™ Burgundy

Capella™ Cherry Vanilla

Capella™ Coral

Capella™ Fuchsia Lace

Capella™ Hello Yellow

Capella™ Indigo Imp.

Capella™ Magenta Diamond

Capella™ Mulberry

DANZIGER

The **CAPELLA™** series

Capella™ Neon Pink

Best Seller

Capella™ Pink Lace

Capella™ Pink Morn

Capella™ Purple

Capella™ Purple Veins

Capella™ Rim Fuchsia

Capella™ Rim Raspberry

Capella™ Rose

Capella™ Ruby Red Imp.

Capella™ Salmon Imp.

Capella™ Sangria

Capella™ White Imp.

DANZIGER

CAPELLA™

For additional information and availability: CAPELLA™

Petunia Hybrida



Stick on priority – Number 2 out of 4 categories

PROPAGATION TIPS

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 72°-74° F (22° - 23°C)	No	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 65° - 68° (18° - 20° C)		Weeks 3-4 100 to 150 ppm	Day of sticking & Week 2

- **Irrigation specification –** Reduce mist as soon as possible

The series is well branched, so pinching is not required for a quality finish. Daminozide at 1500 ppm the day after sticking. Week 4 – Daminozide @ 2000 to 2500 ppm

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.6 to 6.0	0.80	72°-74° F (22° - 23°C)	50 ppm N in mist	Additional lighting is not required	Daminozide day after sticking	Day of sticking	
Week 2	5.6 to 6.0	.80 to .90	72°-74° F (22° - 23°C)	50 ppm N in mist			Second fungicide app	
Week 3	5.6 to 6.0	.90 to 1.0	Cool to 68°F (20°C)	Feed 100 ppm to 150 ppm				
Week 4	5.6 to 6.0	1.0 to 1.2	65° to 68°F (18° to 20°C)	Feed 100 ppm to 150 ppm		Daminozide spray application @ 2000 to 2500 ppm		

DANZIGER

CAPELLA™

For additional information and availability: CAPELLA™

Petunia Hybrida

FINISHING TIPS

- Maintain lower pH levels for optimal Iron (Fe) and Manganese (Mn) availability to plants
- Cool temperature is best for high quality finishing

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks	Average Day	No pinch necessary Additional lighting is not required	150 - 200 ppm N	In liner and finish stages. Daminozide @ 2000 to 2500 ppm
(12 cm) 5” pot	65° to 68° F (18° to 20° C)			
8 to 9 wks	1 plant per pot		Soil EC 1.2 - 1.5 pH 5.6 to 6.0	
(15 cm) 6 ½”				
8 to 10 wks				
(20 cm) 8”	3 plants per container			
8 to 10 wks				
(25 cm) 10”	3 plants per container			

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID,DICHLORVOS
	Thrip	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Transplant to Week 4	5.6 to 6.0	1.2 to 1.5	65° to 68° F (18° to 20° C)	feed 150 to 200 ppm	N/A	Daminozide spray application	Spray fungicide after transplant	
Week 5 to 10	5.6 to 6.0	1.2 to 1.5	65° to 68° F (18° to 20° C)	feed 150 to 200 ppm	N/A	Paclobutrazol Drench at 1 to 2 ppm		Paclobutrazol 1 to 2 ppm drench when plants reach 85% of desired size

- Additional Iron (Fe) or Manganese (Mn) may be required depending on water quality and alkalinity
- PGR - Spray Daminozide 2,000 to 2,500 ppm and / or Paclobutrazol drench application at 1 to 2 ppm for finishing if required

DANZIGER