

# SALLYROSA™ April Night

*Salvia nemorosa*

- First-Year flowering with no bulking or vernalization requirements
- Early and uniform bloom
- Attractive foliage
- Well branched, strong and radial body
- Treat just like an annual



Spring /  
Summer  
Flowering



30 - 41 cm  
12 - 14 in



36 - 41 cm  
14 - 16 in



Zone  
4-9

1<sup>st</sup>  
YEAR  
Flowering



# SALLYROSA™ April Night

For additional information and availability: [Sallyrosa™ April Night](#)



*Salvia nemorosa*

Stick on priority – Number 3 out of 4 categories

PROPAGATION TIPS

Average Time	Temperature	Hormone	Fertilization	Fungicide
4 weeks	Weeks 1 - 2 21° - 22°C (70°-72° F)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 4 20° - 21° C (68° - 70° F)		Weeks 3-4 100 to 125 ppm N	Day of sticking & Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 / 6.2	0.80	21° - 22°C (70°-72° F)	50 ppm N in mist			Day of sticking	
Week 2	5.8 to 6.2	.80 to .90	21° - 22°C (70°-72° F)	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	Cool to 20° - 21° C (68° - 70° F)	Feed 100 ppm to 150 ppm		Daminozide as needed		
Week 4	5.8 to 6.2	1.0 to 1.2	20° - 21° C (68° - 70° F)	feed 100 to 150 ppm	None required	Daminozide as needed		Pinch - Recommended
Week 5	5.8 / 6.2	0.80	20° - 21° C (68° - 70° F)	50 ppm N in mist				



– **Irrigation specification** – Spray with adjuvant the day of sticking/Remove from mist as soon as possible



– *Salvia nemorosa* Sallyrosa™ are day neutral. K-IBA spray application will hasten and even rooting response. Pinch in liner stage or at transplant to increase branching.



# Salvia nemorosa

## Salvia nemorosa

- Cool finishing temperatures will increase quality and longevity of flowers

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
8 to 9 wks	Average Day	Pinch –Optional	150 - 200 ppm N	Daminozide spray application @ 2000 – 3750 ppm Paclobutrazol 1 to 2 ppm drench
10 cm (quart) 1 ppp	(16° to 18°C)			
9 to 11			Soil EC 1.2 – 1.5 pH 5.8 to 6.3	
15 cm (1 gallon) 1 ppp				

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 5 Transplant</b>	5.8 to 6.3	1.0 to 1.2	16° to 18°C	feed 100 to 150 ppm			Drench fungicide after transplant	Control of Pythium & Botrytis
<b>Week 6 to 13</b>	5.8 to 6.3	1.2 to 1.5	16° to 18°C	feed 150 to 200 ppm		Dark Matter may require additional applications		Daminozide spray application @ 2000 – 3750 ppm Paclobutrazol 1 to 2 ppm drench



- PGR - Spray Uniconazole 5 to 7 ppm early in crop  
Paclobutrazol drench application at 0.5 ppm for finishing
- Spray broad spectrum fungicide after transplant
- Scout for Thrip on regular basis

<b>Pests</b>	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrips	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
<b>Diseases</b>	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRDIONE, POLYOXIN
	Pythium Phytophthora	PROPAMOCARB / MEFENOXAM
	Rhizoctonia	AZOXYSTRONBIN / ETRIDIAZOLE / FLUDIOXONIL / PCNB