


BALI™ Blue

Scaevola aemula


- Early flowering variety- About 7 days earlier than touch
- Naturally semi trailing habit
- Abundance of flowers
- Season extender
- Great summer item
- Well suited for hanging baskets and combinations




Semi trailing




20-25 cm
8-10 in



25-41 cm
10-16 in



Heat
tolerant



12-15
Qt, Gal




Landscape



Hanging
basket



Scaevola aemula



Stick on priority – Number 4 out of 4 categories

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 3 72°-74° F (22° - 23°C)	Recommended	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 4 - 5 68° - 70° (20° - 21° C)		Weeks 3-5 100-150 ppm N	Apply broad spectrum Week 2

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8/6.2	0.80	72°-74° F (22° - 23°C)	50 ppm N in mist	Not required			Day of sticking
Week 2	5.8 to 6.2	.80 to .90	72°-74° F (22° - 23°C)	50 ppm N in mist				Second fungicide app
Week 3	5.8 to 6.2	.90 to 1.0	72°-74° F (22° - 23°C)	Feed 100 ppm to 150 ppm				
Week 4	5.8 to 6.2	1.0 to 1.2	68° - 70° (20° - 21° C)	Feed 100 ppm to 150 ppm				
Week 5	5.8 to 6.2	1.0 to 1.2	68° - 70° (20° - 21° C)	Feed 100 ppm to 150 ppm		Daminozide/ uniconzolel as needed		Daminozide @ 2000 to 2500 ppm



- **Misting specification** - Remove from mist as soon as possible
- Do not over-mist or saturate soil - Spray adjuvant at sticking



- Maintain warm temperatures allowing plants to develop.
- Spray Daminozide at 2000 to 2500 ppm or apply Uniconzole 2-5 ppm if required
- Pinch is optional recommended week 4 or at transplant



Scaevola aemula

- Keep the temperatures moderately warm to allow plants to continue to have a good development (average daily temperature at 68 to 70 F (20 to 21 C).

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
6 to 7 wks (12 cm) 5" pot	Average Day 68°-70° F (20°-21° C)	pinch for better branching and uniformity - suggested Night interruption or extended day-Optional	100-150 ppm N Soil EC 1.0 - 1.2 pH 5.8 to 6.2	Spray Daminozide @ 2500 to 3750 ppm Paclobutrazol drench @ 1 to 2 ppm
8 to 9 wks (15 cm) 6 ½"	1 plant per pot			
8 to 9 wks				
(20 cm) 8 "	3 plants per container			

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrip	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 5 Transplant	5.8 to 6.2	1.0 to 1.2	68° to 70° F (20° to 21° C)	feed 100 to 150 ppm		B-9 as needed	Spray fungicide after transplant	Spray Daminozide @ 2500 to 3750 ppm
Week 6 to 13	5.8 to 6.2	1.0 to 1.2	68° to 70° F (20° to 21° C)	feed 100 to 150 ppm		Uniconzolel if required		Paclobutrazol drench @ 1 to 2 ppm



- **PGR** - Spray Daminozide at 2500 to 3750 ppm for finishing plants
- Uniconzole 2-5 ppm if required for finishing.
- **Drench** a broad spectrum fungicide after transplant
- **Scout** for Aphids and Fungus Gnats and Mites